

Cardinality revisited

- A set is finite (has finite cardinality) if its cardinality is some (finite) integer n .
- Two sets A, B have the same cardinality iff there is a one-to-one correspondence from A to B
- E.g. alphabet (lower case)
- a b c
- 1 2 3

Infinite sets

- Why do we care?
- Cardinality of infinite sets
- Do all infinite sets have the same cardinality?

Countable sets

Defn: Is finite OR has the same cardinality as the positive integers.

- Why do we care?

E.g.

- The algorithm works for “any n ”
- Induction!

A special countable set

- The set of all binary strings
- Therefore the set of all Java programs is countable!

Countable sets – contd.

- Proving this involves (usually) constructing an explicit bijection with positive integers.
- Fact (Will not prove): Any subset of a countable set is countable.

Will prove that

- The rationals are countable!
- The reals are not countable

Countably Infinite Sets

A set S is infinite if there exists a surjective function $F: S \rightarrow \mathbb{N}$.

“The set \mathbb{N} has no more elements than S .”

A set S is countable if there exists a surjective function $F: \mathbb{N} \rightarrow S$

“The set S has not more elements than \mathbb{N} .”

A set S is countably infinite if there exists a bijective function $F: \mathbb{N} \rightarrow S$.

“The sets \mathbb{N} and S are of equal size.”

The integers are countable

- Write them as
0, 1, -1, 2, -2, 3, -3, 4, -4,
- Find a bijection between this sequence
and 1,2,3,4,.....

Notice the pattern:

$1 \rightarrow 0$	$2 \rightarrow 1$	So $f(n) = n/2$ if n even $-(n-1)/2$ o.w.
$3 \rightarrow -1$	$4 \rightarrow 2$	
$5 \rightarrow -2$	$6 \rightarrow 3$	

Other simple bijections

- Odd positive integers

$$1 \rightarrow 1 \quad 2 \rightarrow 3 \quad 3 \rightarrow 5 \quad 4 \rightarrow 7 \dots$$

- Union of two countable sets A, B is countable:

Say $f: \mathbb{N} \rightarrow A$, $g: \mathbb{N} \rightarrow B$ are bijections

New bijection $h: \mathbb{N} \rightarrow A \cup B$

$$h(n) = f(n/2) \text{ if } n \text{ is even}$$

$$= g((n-1)/2) \text{ if } n \text{ is odd.}$$

The rationals are countable

- Show that $\mathbb{Z}^+ \times \mathbb{Z}^+$ is countable.
- Trivial injection between \mathbb{Q}^+ , $\mathbb{Z}^+ \times \mathbb{Z}^+$.
- To go from \mathbb{Q}^+ to \mathbb{Q} , use the trick used to construct a bijection from \mathbb{Z} to \mathbb{Z}^+ .
- Details on the board.

Facts to note

- Note that the ordering of \mathbb{Q} is not in increasing order or decreasing order of value.
- In proofs, you CANNOT assume that an ordering has to be in increasing or decreasing order.
- So cannot use ideas like “between any two real numbers x , y , there exists a real number $0.5(x+y)$ ” to prove uncountability.

The reals are not countable

- Wrong proof strategy:
 - Suppose it is countable
 - Write them down in increasing order
 - Prove that there is a real number between any two successive reals.
 - WHY is this incorrect?
- (Note that the above “proof” would show that the rationals are not countable!!)

The reals are not countable - 2

- Cantor diagonalization argument (1879)
- **VERY** powerful, important technique.
- Proof by contradiction.
- Strategy
 - Assume countable
 - look at all numbers in the interval $[0,1)$
 - list them in ANY order
 - show that there is some number not listed

Uncountable Sets

There are infinite sets that are not countable.
Typical examples are \mathbb{R} , $\mathcal{P}(\mathbb{N})$ and $\mathcal{P}(\{0,1\}^*)$

We prove this by a diagonalization argument.
In short, if S is countable, then you can make
a
list s_1, s_2, \dots of all elements of S .

Diagonalization shows that given such a list,
there will always be an element x of S that
does not occur in s_1, s_2, \dots

Uncountability of $\mathcal{P}(\mathbb{N})$

The set $\mathcal{P}(\mathbb{N})$ contains all the subsets of $\{1, 2, \dots\}$. Each subset $X \subseteq \mathbb{N}$ can be identified by an infinite string of bits $x_1 x_2 \dots$ such that $x_j = 1$ iff $j \in X$.

There is a bijection between $\mathcal{P}(\mathbb{N})$ and $\{0, 1\}^{\mathbb{N}}$.


Proof by contradiction: Assume $\mathcal{P}(\mathbb{N})$ countable.

Hence there must exist a surjection F from \mathbb{N} to the set of infinite bit strings.

“There is a list of *all* infinite bit strings.”

Diagonalization

Try to list all possible infinite bit strings:



0	0	0	0	0	0	...
1	1	1	1	1	1	...
2	1	0	0	0	0	...
3	0	1	0	1	0	...
⋮						⋮

Look at the bit string on the diagonal of this table: 0101... The negation of this string (“1010...”) does not appear in the table.

No Surjection $N \rightarrow \{0,1\}^N$

Let F be a function $N \rightarrow \{0,1\}^N$.

$F(1), F(2), \dots$ are all infinite bit strings.

Define the infinite string $Y = Y_1 Y_2 \dots$ by

$$Y_j = \text{NOT}(j\text{-th bit of } F(j))$$

On the one hand $Y \in \{0,1\}^N$, but on the other hand:
for every $j \in N$ we know that $F(j) \neq Y$
because $F(j)$ and Y differ in the j -th bit.

F cannot be a surjection: $\{0,1\}^N$ is uncountable.

The set of all functions

-is therefore uncountable!
- So there must exist problems for which there do not exist Java programs (or pseudocode, or algorithms!)

Generalization

- We proved that $P(\{0,1\}^*)$ is uncountably infinite.
- Can be generalized to $P(\Sigma^*)$ for any finite Σ .

R is uncountable

- Similar diagonalization proof. We will prove $[0,1)$ uncountable
- Let F be a function $N \rightarrow R$
 $F(1), F(2), \dots$ are all infinite digit strings (padded with zeroes if required).
- Define the infinite string of digits $Y = Y_1 Y_2 \dots$ by
$$Y_j = \begin{cases} F(i)_i + 1 & \text{if } F(i)_i < 8 \\ 7 & \text{if } F(i)_i \geq 8 \end{cases}$$

Q: Where does this proof fail on N ?

Other infinities

- We proved $2^{\mathbb{N}}$ uncountable. We can show that this set has the same cardinality as $\mathcal{P}(\mathbb{N})$ and \mathbb{R} .
- What if we take $\mathcal{P}(\mathbb{R})$?
- Can we build bigger and bigger infinities this way?
- Cantor: **Continuum hypothesis** – YES!

Notes

- The cardinality of neither the reals nor the integers are finite, yet one set is countable, the other is not.
- Q: Is there a set whose cardinality is “in-between”?
- Q: Is the cardinality of \mathbb{R} the same as that of $[0,1)$?