

EECS 1028 M: Discrete Mathematics for Engineers

Suprakash Datta

Office: LAS 3043

Course page: <http://www.eecs.yorku.ca/course/1028>

Also on Moodle

Administrivia

- Lectures: Mon-Wed-Fri 1:30-2:30 pm (LAC C)
- Tests (35%) : 3 tests, 15% each (worst test to be scaled to 5%)
- final (40%)
- Homework (15%),
- Tutorials(10%)
- Office hours: Mon-Wed 3-4 pm or by appointment at LAS 3043.

Textbook: Kenneth H. Rosen. Discrete Mathematics and Its Applications, 7th Edition. McGraw Hill, 2012.

Homework, Grades

- We will be paperless, except for tests, quizzes, final examination.
- All course information online – Moodle, partly mirrored in public course webpage
- All homework **MUST** be typed. You will get a zero if you submit handwritten solutions. You may use Office, Google Docs, LaTeX, or other packages but all submissions must be in pdf format.
- We will use crowdmark for grading. Follow instructions for re-appraisal requests.
- All returned work will be in electronic form (including quizzes, tests).
- I would like to use iClicker if possible.

Tutorials, Quizzes

- Tutorials (2 hours/week) are **mandatory**. Attendance will be taken.
- You get 0.5% for each tutorial attended.
- Every other week, there will be a short quiz (10-15 min), mirrored closely after the tutorial problems covered in the last 1 or 2 tutorials. Each quiz is worth 2%, but your attendance points are added to it to a maximum of 2%.
- Missed quizzes cannot be made up. If you have a valid medical reason, the weight will be transferred to the final.
- Missed tests also cannot be made up. If you have a valid medical reason, the weight will be transferred to the final.
- If you have serious non-medical reasons (having work is not one), talk to me. We will deal with those on an ad hoc basis.

Course Objectives

We will focus on two major goals:

- Basic tools and techniques in discrete mathematics
 - Set Theory, Functions and Relations
 - Propositional and Predicate logic
 - Sequences and Series
 - Induction, Recursion
 - Simple Combinatorics
 - Introductory Graph Theory
- Precise and Rigorous Mathematical Reasoning
 - Writing proofs

My Expectations

- You will attend classes and tutorials regularly
- Want to solidify your Math foundations
- Ask for help when needed
- Follow academic honesty regulations (see the class webpage for more details on policies).

To do well in this class

- Study with pen and paper
- Ask for help early
- Practice, practice, practice ...
- Follow along in class rather than take notes
- Ask questions in class or outside class
- Keep up with the class
- Read the book, not just the slides
- Be timely – HW submitted late will not be graded

Mathematical Reasoning

- Why Mathematics?
 - Mathematics as a precise language
 - Precision in definitions
 - Precision in statements
- Motivation (for EECS)
 - Specification (description, modeling)
 - Reasoning (Making precise, rigorous claims)
- Procedure
 - Axioms
 - Inference
 - Facts/Theorems

Examples of Reasoning about Problems

- $0.9999999999999999 \dots = 1?$
- There exists integers a, b, c that satisfy the equation $a^2 + b^2 = c^2$
- There exists integers a, b, c that satisfy the equation $a^4 + b^4 = c^4$
- There are as many integers as there are rational numbers
- The program that I wrote never hangs (i.e. always terminates)
- The program that I wrote works correctly for all valid inputs
- There does not exist an algorithm to check if a given program never hangs

Proofs and Similar Structures

- Backing up statements with reasons
- Providing detailed explanations (Amazon, Netflix,...)
- Understanding the basis of eCommerce, Bitcoin, ...
- Evidence in data mining systems

Intuitive Proofs

- What?
- Why?
- When?
- How much detail?

Review of Fundamentals

- Sets
- Number Systems
- Basic algebra

Set Theory Fundamentals

- Unordered collection of elements, e.g.,
 - Single digit integers
 - Non-negative integers
 - Faces of a die
 - Sides of a coin
 - Students enrolled in EECS1028, W 2018
- Two key aspects of sets:
 - No duplicates
 - No inherent ordering of elements

Set Theory Fundamentals - 2

- $\mathbb{N}$: the set of natural numbers, $\mathbb{R}$: the set of real numbers
- Membership
Notation: $a \in A, b \notin A$
- Ordered pairs
Notation: $(a, b), a \in A, b \in B$
- Equality of sets

Describing Sets

- English description
 - The set of natural numbers between 5 and 8 (inclusive).
 - The set of all students enrolled in EECS1028 M, W 2018
- Enumeration
 - $S = \{1, 2, 3\}$
 - $S = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$
- Set builder-notation
 - $S = \{x \in \mathbb{N} \mid x > 3\}$
 - $S = \{(x, y) \mid x, y \in \{0, 1\}\}$

More on Sets

Special sets

- Universal set U
- Empty set ϕ (How many elements?)

Sets vs Sets of sets

- $\{1, 2\}$ vs $\{\{1, \}, \{2\}\}$
- $\{\}$ vs $\{\{\}\} = \{\phi\}$



Note:

- 1 Connection with data types (e.g., in Java)
- 2 The elements of a set can be sets, pairs of elements, pairs of pairs, triples, ... !!

Set Operations

- Subsets: $A \subseteq B : \forall x(\text{ if } x \in A \text{ then } x \in B)$
- Union: $A \cup B = \{x | (x \in A) \text{ or } (x \in B)\}$
- Intersection: $A \cap B = \{x | (x \in A) \text{ and } (x \in B)\}$
- Difference: $A - B = \{x | (x \in A) \text{ and } (x \notin B)\}$
- Complement: $A^c \text{ or } \bar{A} = \{x | x \notin A\} = U - A$
- Cartesian product: $A \times B = \{(a, b) | a \in A, b \in B\}$
 - "Set of ordered pairs"
 - $\mathbb{R} \times \mathbb{R} = \{(x, y) | x \in \mathbb{R}, y \in \mathbb{R}\}$, "Coordinate plane" or "the real plane" Sometimes called $\mathbb{R}^2$.

Venn Diagrams



More on Sets

- Cardinality - number of (distinct) elements
- Finite set - cardinality some finite integer n
- Infinite set - a set that is not finite
- Power set - Set of all subsets; Notation $\mathcal{P}(S) = \{A | A \subseteq S\}$, sometimes written 2^S

Laws on Set operations

Page 130

- Associative laws

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

- Distributive laws:

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$


- De Morgan's laws

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

Proofs of Laws of Set Operations

Proofs can be done with Venn diagrams.



Number Systems

- Natural numbers, $\mathbb{N} : \{1, 2, 3, \dots\}$.
- Whole numbers, $\mathbb{W} : \{0, 1, 2, 3, \dots\}$.
- Integers, $\mathbb{Z} : \{\dots, -2, -1, 0, 1, 2, 3, \dots\}$
Notation: $\mathbb{Z}^+ = \text{positive integers} = \mathbb{N}$
- Real numbers, $\mathbb{R}$
Notation: $\mathbb{R}^+ = \text{positive reals}$
- Complex numbers, $\mathbb{C} = \{x + iy | x, y \in \mathbb{R}, i^2 = -1\}$.
- Co-ordinates on the plane, $\mathbb{R}^2$.
- Rational numbers, $\mathbb{Q} = \{\frac{m}{n} | m, n \in \mathbb{Z}, n \neq 0\}$.
- Irrational numbers, $\mathbb{R} - \mathbb{Q}$: all real numbers that are not rational. Examples: $\pi, e, \sqrt{2}$.

Number Systems - Questions

- How do we know π , e , $\sqrt{2}$ are not rational?
- How are real numbers represented on a computer?
- Do all rational numbers have finite decimal representations?
Counterexample: $1/3$
- If a number has an infinite decimal representation, can we conclude it is irrational?

Basic Algebra

- Operations with exponents: Theorem 1, pg A-7
 - $b^x * b^y = b^{x+y}$
 - $b^x / b^y = b^{x-y}$
 - $(b^x)^y = b^{xy}$
- Logarithms: Theorem 2, pg A-8
 - Logarithm of products, powers
 - Change of bases
- Operations with polynomials
- Solving linear and quadratic equations