

## Functional Dependencies

Suppose we have a relation  $R$  with attributes  $ABCD$

### 1. What an FD means.

Suppose the functional dependency  $BC \rightarrow D$  holds in  $R$ . Create an instance of  $R$  that violates this FD.

#### Solution:

In order to violate this FD, we need two tuples with the same value for  $B$  and  $C$ , yet different values for  $D$ .

A	B	C	D
1	3	6	4
2	3	6	5

### 2. Equivalent sets of FDs.

(a) Are the sets  $A \rightarrow BC$  and  $A \rightarrow B, A \rightarrow C$  equivalent? If yes, explain why. If no, construct an instance of  $R$  that satisfies one set of FDs but not the other.

#### Solution:

These are equivalent – there is no instance of the relation that satisfies one but not the other. This can be proven, and now that you know the closure test, the proof is very concise:

- Assume that  $A \rightarrow BC$ .
  - Under this assumption,  $A^+ = ABC$ .
  - Therefore  $A \rightarrow B$ , and  $A \rightarrow C$ .
- Assume that  $A \rightarrow B$ , and  $A \rightarrow C$ .
  - Under this assumption,  $A^+ = ABC$ .
  - Therefore  $A \rightarrow BC$ .
- Therefore each set of FDs follows from the other. They are equivalent.

In fact we can always “split the RHS” of an FD. Review the definition of an FD to see why this makes sense.

(b) Are the sets  $AB \rightarrow C$  and  $A \rightarrow C, B \rightarrow C$  equivalent? If yes, explain why. If no, construct an instance of  $R$  that satisfies one set of FDs but not the other.

#### Solution:

These are *not* equivalent, as demonstrated by the following instance. It satisfies  $AB \rightarrow C$  but not  $A \rightarrow C, B \rightarrow C$ :

A	B	C
1	2	4
3	2	5

We can never split the LHS of an FD. Again, the definition of FD makes clear why.

(c) Are the sets  $AB \rightarrow C$  and  $A \rightarrow B, A \rightarrow C$  equivalent? If yes, explain why. If no, construct an instance of  $R$  that satisfies one set of FDs but not the other.

#### Solution:

These are *not* equivalent, as demonstrated by the following instance that satisfies  $AB \rightarrow C$  but not  $A \rightarrow B, A \rightarrow C$ :

A	B	C
1	2	4
1	8	9

### 3. Keys and FDs.

(a) We claimed that if a set of attributes  $K$  functionally determines all attributes,  $K$  must be a superkey (i.e., no two tuples can agree on all attributes in  $K$ ). Do you believe this? Suppose these FDs hold in  $R$ :  $A \rightarrow BC$ ,  $C \rightarrow D$ . Does  $A$  functionally determine all attributes of  $R$ ? Can two tuples agree on  $A$ ?

**Solution:**

This is left as an exercise to explore on your own, in order to build your intuition.

(b) We also said that if  $K$  is a superkey (i.e., no two tuples can agree on all attributes in  $K$ )  $K$  must functionally determine all attributes. Do you believe this? Suppose  $A$  is a superkey of  $R$ . Does  $A$  functionally determine all attributes of  $R$ ?

**Solution:**

Again, this is left as an intuition-building exercise.

### 4. Does an FD follow from a set of FDs?

Suppose we have a relation on attributes  $ABCDEF$  with these FDs:  $AC \rightarrow F$ ,  $CEF \rightarrow B$ ,  $C \rightarrow D$ ,  $DC \rightarrow A$ .

(a) Does it follow that  $C \rightarrow F$ ?

(b) Does it follow that  $ACD \rightarrow B$ ?

**Solution:**

We use the closure test to check whether an FD follows from a set of FDs.

$C^+ = CDAF$ . Therefore,  $C \rightarrow F$  does follow.

$ACD^+ = ACDF$ . Therefore,  $ACD \rightarrow B$  does not follow.

### 5. Projecting a set of FDs onto a subset of the attributes.

Suppose we have a relation on attributes  $ABCDE$  with these FDs:

$$A \rightarrow C, C \rightarrow E, E \rightarrow BD$$

(a) Project the FDs onto attributes  $ABC$ .

**Solution:**

To project onto a set of attributes, we systematically consider every possible LHS of an FD that might hold on those attributes.

- $A^+ = ACEBD$ , therefore  $A \rightarrow BC$ . (It also functionally determines  $DE$ , but these are not in our set of attributes. And it functionally determines itself, but we don't need to write down dependencies that are tautologies.)
- $B^+ = B$ . This yields no FDs for our set of attributes.
- $C^+ = CEBD$ , therefore  $C \rightarrow B$ .
- We don't need to consider any supersets of  $A$ .  $A$  already determines all of our attributes  $ABC$ , so supersets of  $A$  will only yield FDs that already follow from  $A \rightarrow BC$ .
- The only remaining subset of the attributes  $ABC$  to consider is  $BC$ .  $BC^+ = BCED$ . This yields no FDs for our set of attributes.
- So the projection of the FDs onto  $ABC$  is:  $\{A \rightarrow BC, C \rightarrow B\}$ .

(b) Project the FDs onto attributes  $ADE$ .

**Solution:**

- $A^+ = ACEBD$ , therefore  $A \rightarrow DE$ .
- $D^+ = D$ . This yields no non-trivial FDs.
- $E^+ = EBD$ , therefore  $E \rightarrow D$ .
- Again, we don't need to consider any supersets of A, since A determines all the attributes ADE also.
- The only remaining subset of the attributes  $ABC$  to consider is  $DE$ .  $DE^+ = DEB$ . This yields no FDs for our set of attributes.
- So the projection of the FDs onto  $ADE$  is:  $\{A \rightarrow DE, E \rightarrow D\}$ .